

## FLARAXIN APPLICATION EFFICIENCY IN DIFFERENT CASES WITH CANCER.

Scientific and Treatment Center for Cancer Curing PHOENIX, Kharkov, Ukraine

Dr.Elena Oleynik, Dr.Tatyana Parshina, Dr. Galina Schukina

<b>Application efficiency of FLARAXIN at different oncological diseases.</b>			
Maximum, (+ + +)	High, (+ +)	Average, (+)	Non-efficient, (-)
<ol style="list-style-type: none"> <li>1. Melanoma;</li> <li>2. Tumors of the mammary gland;</li> <li>3. Tumor of lungs, little worse bronchi tumors;</li> <li>4. Cancer of the small intestine;</li> <li>5. Cancer of the colon;</li> <li>6. Cancer of the rectum;</li> <li>7. Cancer of the prostate gland;</li> <li>8. Non-malignant tumors;</li> <li>9. Adenoma;</li> <li>10. The tumors of feminine sexual organs:               <ol style="list-style-type: none"> <li>a) Fibromioma;</li> <li>б) Cervix of the uterus Cancer;</li> <li>в) Cancer of womb body;</li> <li>г) non-malignant tumors of gonads, polycystic (calcined);</li> <li>д) Cancer of gonads;</li> </ol> </li> <li>11. Tumors of CNS (Tumors of Cerebrum (Brain)):               <ol style="list-style-type: none"> <li>A) Ependioma – non-malignant;</li> <li>Б) Oligodendroglioma;</li> <li>B) Astrocitoma;</li> <li>Г) Tumors of pituitary gland;                   <ol style="list-style-type: none"> <li>Г1) Adenoma of pituitary gland;</li> <li>Г2) Carnofaringioma;</li> </ol> </li> <li>Д) Brain metastases.</li> </ol> </li> </ol>	<ol style="list-style-type: none"> <li>1. Neoplasms of maxillofacial areas:               <ol style="list-style-type: none"> <li>Cancer of the lips, tongue cancer;</li> </ol> </li> <li>2. Tumor of the upper respiratory tract and bodies;</li> <li>3. Cancer of the maxillary sinus;</li> <li>4. Tumor of the pharynx;</li> <li>5. Tumor of the larynx;</li> <li>6. Mediastinal tumors;</li> <li>7. Tumors of the stomach;</li> <li>8. Pancreatic cancer;</li> <li>9. Tumors of the kidney: adenocarcinoma;</li> <li>10. Bladder cancer.</li> </ol>	<ol style="list-style-type: none"> <li>1. Sarcoma of soft tissues of any etiology;</li> <li>2. Sarcoma of bone (depends on the blood supply to tumors);</li> <li>3. Tumors of the esophagus (at the last stage IV could only prolong life, but in the initial stages till III stage – can be treated well);</li> <li>4. Metastases in Liver (liver cancer - is inefficient, but metastases to the liver - can be well treated);</li> <li>5. Tumors of the testis;</li> <li>6. Tumors of the penis.</li> </ol>	<ol style="list-style-type: none"> <li>1. Diseases of blood, leukemia, lymphogranulomatosis (immunosuppression should be done, and vice versa Flaraxin raises immunity);</li> <li>2. Extensive liver disease;</li> <li>3. Intoxication (moderate, high) are required to reduce the level of intoxication, because flaraxin destroying tumor tissue and by this way increases the level of intoxication;</li> </ol> <p style="text-align: center;">*</p>

\* If you have high pain level. The work of flaraxin is complicated and it can give more pain in the first days of application.